

FREQUENCY OF OCCURRENCE AND VEGETATION SELECTION BY SOUTH TEXAS

BATS

A Thesis

by

KYLIE PEREZ

Submitted to the Graduate and Professional School of  
Texas A&M University  
in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE

|                     |                 |
|---------------------|-----------------|
| Chair of Committee, | Roel R. Lopez   |
| Committee Members,  | Andrea Bruno    |
|                     | David M. Cairns |
| Head of Department, | Roel R. Lopez   |

August 2023

Major Subject: Wildlife and Fisheries Sciences

Copyright 2023 Kylie F. Perez

## ABSTRACT

Bats are an ecologically diverse order and important contributors to natural ecosystems. Despite their importance, detailed information on the ecology of bats is limited within the southern portion of Texas where most species found throughout the United States are located. My objective was to provide a basic understanding of the frequency of occurrence and vegetation preference by South Texas bats to aid in their management and conservation. I monitored 7 resident bat species: Evening bat (*Nycticeius humeralis*), Eastern red bat (*Lasiurus borealis*), Hoary bat (*Lasiurus cinereus*), Northern yellow bat (*Lasiurus intermedius*), Mexican free-tailed bat (*Tadarida brasiliensis*), Tri-colored bat (*Perimyotis subflavus*), and the Cave myotis bat (*Myotis velifer*) in the Texas Coastal Plains of South Texas at the El Sauz Ranch (East Foundation) throughout February-July 2022. Acoustic data were collected using acoustic monitors and the data for 25 unique nights at 25 distinct sampling locations, within 3 vegetation groups were obtained. The three vegetation groups used in my study included woodland/shrubland, rangeland/wetland, and sand. A total of 20,518 sound files were used in my analysis revealing use showed a preference for the woodland/shrubland composition with a vegetation-selection ratio of  $S=4.56$  as well as the sand composition ( $S=3.01$ ), and avoidance for the rangeland/wetland composition ( $S=0.35$ ). Management practices should focus on landscape heterogeneity with a notation on the importance of conserving native woodland/shrubland areas.

## DEDICATION

This thesis is dedicated to the people who have supported me through my education. To my parents, Rudy and Kelli Perez, for their unwavering support, thank you for helping me see this journey through to the end.

## ACKNOWLEDGMENTS

I would like to thank my committee chair, Dr. Roel Lopez for providing guidance and funding for my research and education as well as Dr. Michael Morrison for project guidance and support. I appreciate them for believing in me as a student, a scientist, and a person.

I would like to thank my committee members, Dr. Andrea Montalvo and Dr. David Cairns, for their project support as well. During my initial wildlife introduction to South Texas, Dr. Montalvo played a key role in instilling a passion for this field and developing leadership skills. I also would like to thank the East Foundation for allowing me access to their ranches, accommodations, and support through ranching, science, and education to promote the advancement of land stewardship. I would like to express my gratitude to my lab mates and colleagues for their invaluable guidance in my academic pursuits and personal growth. Their dedication to hard work and enjoyment of this time in our lives has been an inspiration.

Lastly, I would like to thank my family. Especially my parents, Rudy and Kelli Perez, for supporting me 100% in anything I wish to do. Thank you for your generosity in helping me through college and life and instilling in me a passion for animals and curiosity. They inspire me to constantly strive towards the best version of myself.

## CONTRIBUTORS AND FUNDING SOURCES

### **Contributors**

This work was supported by a thesis committee consisting of Dr. Roel Lopez of the Department of Rangeland, Wildlife and Fisheries Management, Dr. Andrea Montalvo of the East Foundation, and Dr. David Cairns of the Department of Geography. All data collected for the thesis were completed by the student independently.

### **Funding Sources**

The graduate study was supported by a Graduate Assistantship from Dr. Michael Morrison and Texas A&M University.

## NOMENCLATURE

|      |                         |
|------|-------------------------|
| ESR  | El Sauz Ranch           |
| kHz  | Kilo Hertz              |
| m    | Meter                   |
| km   | Kilometer               |
| Labo | Eastern red bat         |
| Laci | Hoary bat               |
| Lain | Northern yellow bat     |
| Myve | Cave myotis bat         |
| Nyhu | Evening bat             |
| Pesu | Tri-colored bat         |
| Tabr | Mexican free-tailed bat |

TABLE OF CONTENTS

|                                       | Page |
|---------------------------------------|------|
| ABSTRACT.....                         | ii   |
| DEDICATION.....                       | iii  |
| ACKNOWLEDGMENTS .....                 | iv   |
| CONTRIBUTORS AND FUNDING SOURCES..... | v    |
| NOMENCLATURE .....                    | vi   |
| TABLE OF CONTENTS .....               | vii  |
| LIST OF FIGURES .....                 | ix   |
| LIST OF TABLES .....                  | xi   |
| INTRODUCTION .....                    | 1    |
| METHODS .....                         | 4    |
| Study Area .....                      | 4    |
| Sampling and Data Collection .....    | 4    |
| DATA ANALYSIS.....                    | 8    |
| Acoustic Data Analysis.....           | 8    |
| Vegetation Selection Ratios .....     | 9    |
| RESULTS .....                         | 11   |

|                                   |        |
|-----------------------------------|--------|
| Frequency of Occurrence .....     | 11     |
| Vegetation Selection .....        | 11     |
| Eastern red bat.....              | 16     |
| Hoary bat.....                    | 17     |
| Northern yellow bat .....         | 18     |
| Cave myotis bat.....              | 19     |
| Evening bat .....                 | 20     |
| Tri-colored bat .....             | 21     |
| Mexican free-tailed bat .....     | 22     |
| <br>DISCUSSION.....               | <br>23 |
| <br>MANAGEMENT IMPLICATIONS ..... | <br>27 |
| <br>REFEERENCES .....             | <br>28 |

## LIST OF FIGURES

|   | Page |
|---|------|
| Figure 1. Sites on the ESR where acoustic data were collected for the following vegetation compositions: woodland/shrubland, rangeland/wetland, and sand.....   | 7    |
| Figure 2. Sonobat call analysis sonogram.....   | 9    |
| Figure 3. Vegetation availability by composition; rangeland/wetland (82%), woodland/shrubland (12%), sand (5%), other (1%).....   | 12   |
| Figure 4. Total vegetation-selection ratios ( $S$ ) ( $S > 1$ signifies a preference for the area, $S < 1$ signifies an avoidance) for each vegetation composition, woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....   | 14   |
| Figure 5. Species specific selection ratios ( $S$ ) for each vegetation composition ( $S > 1$ signifies a preference for the area, $S < 1$ signifies an avoidance), Eastern red bat (Labo), Hoary bat (Laci), Northern yellow bat (Lain), Cave myotis bat (Myve), Evening bat (Nyhu), Tri-colored bat (Pesu), Mexican free-tailed bat (Tabr), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD)..... | 15   |
| Figure 6. Eastern red bat vegetation-selection ratios stratified by vegetation type ( $S > 1$ signifies a preference for the area, $S < 1$ signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....   | 16   |
| Figure 7. Hoary bat vegetation-selection ratios stratified by vegetation type ( $S > 1$ signifies a preference for the area, $S < 1$ signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....   | 17   |
| Figure 8. Northern yellow bat vegetation-selection ratios stratified by vegetation type ( $S > 1$ signifies a preference for the area, $S < 1$ signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....   | 18   |

Figure 9. Cave myotis bat vegetation-selection ratios stratified by vegetation type ( $S > 1$  signifies a preference for the area,  $S < 1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....19

Figure 10. Evening bat vegetation-selection ratios stratified by vegetation type ( $S > 1$  signifies a preference for the area,  $S < 1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....20

Figure 11. Tri-colored bat vegetation-selection ratios stratified by vegetation type ( $S > 1$  signifies a preference for the area,  $S < 1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....21

Figure 12. Mexican free-tailed bat vegetation-selection ratios stratified by vegetation type ( $S > 1$  signifies a preference for the area,  $S < 1$  signifies an avoidance), woodland/shrubland WS), rangeland/wetland (RW), and sand (SD).....22

## LIST OF TABLES

|  | Page |
|--|------|
| Table 1. Scientific and common names for every species analyzed with total occurrence per species for the entirety of the study in decreasing order of occurrence..... | 11   |
| Table 2. Studied bat species with total occurrence per vegetation composition, woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).....                     | 13   |

## INTRODUCTION

Despite being more ecologically diverse than any other group of mammals and vital members of the natural ecosystems, many aspects of bats such as fine-temporal resource selection are widely understudied in “hot zones” throughout the southern portion of the United States. Bats belong to the 2nd largest order of mammals called Chiroptera (Bernard 2002). These highly innovative animals have developed a multitude of adaptations enabling them to navigate and exploit diverse ecological niches, resources, and habitats. Echolocation is one of these refined adaptations, a characteristic that is only prevalent in the bat suborder Microchiroptera and dolphins (Schnitzler and Kalko 2001). Animals use echolocation to locate objects by emitting high-frequency signals and interpreting the reflected echoes (Schnitzler and Kalko 2001). This method is employed by bats during the search for insects and agricultural pests as prey. When studying foraging behavior, the majority of research has relied on ultrasonic detectors to monitor echolocation calls, rather than directly measuring the activity of individual animals (Russo and Jones 2002).

Bats provide extremely important ecosystem services, like insect control, in regulating and improving rangelands with growing crops and livestock (Kunz et al. 2011). Bats feed on mosquitoes and other pests that are crucial to control in order to conserve land. Results of an original study concluded that a colony of bats can consume nearly two-thirds of its body weight in insects every night (Coutts et al. 1973). Some bat species are also important pollinators by feeding and dispersing seeds for nearly 300 species of fruits (Sekercioglu 2006). Bats also serve as food for owls, hawks, and raccoons. Consumptive goods indicating economic value provided by bats such as bat waste, called guano, can be used as fertilizer for agricultural plots and home

gardens as it is composed of rich nutrients including nitrogen, phosphorus, and potassium (Kunz et al. 2011, Campbell 2022).

Birds are often cited in the literature as important bioindicators in agriculture, but a study done in 2008 showed insectivorous bats have stronger ecological effects than birds (Kalka et al. 2008). Extended knowledge of bat practices is crucial to determine the most appropriate methods of conservation. This study, focusing on species that reside in South Texas specifically, would provide much-needed information to aid in future conservation and management efforts.

Texas is home to almost 70% of the species of bats found in the United States (Knight 2020). However, Texas bats receive only a small fraction of scholarly attention, with only around 3% of publications focus on specific species (Fern et al. 2018). Texas also possesses the largest known bat colony in the world, Bracken Cave Preserve, which is also one of the largest concentrations of mammals on earth (Woodward 2022). Among other reasons, Texas is an epicenter for bats and therefore provides an abundance of opportunities for research to be done.

Currently, pesticides are the primary means of controlling agricultural pests, which can place wildlife at risk of chemical exposure (Agosta 2002). This may be a factor for decline in bat population numbers. Pesticides may cause coordination loss, weaken animals, and compromise flight and the ability to orient (Torquetti et al. 2021). In promoting environmental sustainability and supporting the conservation of at-risk species (i.e., bats), contemporary consumers show a growing preference for agri-food products (Russo et al. 2018). The use of higher amounts of fertilizer and agrochemical use generally precedes a reduction in biodiversity (Grashof-Bokdam and Van Langevelde 2005, Billeter et al. 2008, Liira et al. 2008). This is consistent with an established ecological theory that suggests an inverse correlation between site productivity and

species diversity i.e., increased productivity leads to a decrease in diversity (Gough et al. 2000, Grime 2001, Mittelbach et al. 2001, Liira et al. 2008).

Bat-habitat relationships are poorly understood because of the high mobility of bats, which gives them access to a wide range of territories. A comprehensive study of how bat species interact with their environment should be the driving factor in the management of the species, community, or ecosystem. The objective of my study is to provide a baseline for the frequency of occurrence and vegetation selection by bats establishing South Texas as their environment and lead to future, specific studies. Based on vegetation availability, I would expect to see the most total frequency of occurrence for all species within the rangeland/wetland composition as this vegetation is the most available as well as providing ample area for foraging (Morris et al. 2011, Barros et al. 2014, Trubitt et al. 2019, Wied et al. 2020). This information will provide ranchers with a better understanding of the land they manage and how to maintain biodiversity for sustainable range management. As a result, ranchers will be proactive in sustaining the ecological robustness of their lands.

## METHODS

### Study Area

This study was conducted at the El Sauz Ranch (ESR), belonging to the East Foundation, near Port Mansfield, TX in the Texas Coastal Plains (Texas Parks and Wildlife Department) in Willacy County (26.536769° N, -97.446649° W). The East Foundation was recently named the nation's first Agriculture Research Organization (ARO) in early 2022 (Marin 2023). Their mission is to promote the advancement of land stewardship through ranching, science, and education. The mean monthly temperature of the study area during the study period (February-July 2022) was 25.4 C with a mean minimum temperature of 13.2 C and a mean maximum temperature of 37.7 C (PRISM Climate Group 2023). The mean precipitation of the study area during the study period was 60.9 mm (PRISM Climate Group 2023). The study area used for this study spans 10,981.72 hectares with a variety of unique vegetation compositions. The ESR is composed of woodland/shrubland (1,380 ha), rangeland/wetland (8,982 ha), and sand/bare ground (523 ha). Mostly mid to tall grasses (cordgrass [*Spartina alterniflora*]) as well as prickly pear (*Opuntia And*) and catclaw (*Uncaria tomentosa*) cover this ecoregion, with an overstory of honey mesquite (*Prosopis glandulosa*) and live oak (*Quercus virginiana*) trees (Cooper et al. 2021). This area is also home to mostly stabilized and active sand dune deposits. The ESR is a working cattle ranch operation and home to multiple research programs. I selected this area for my research as it is representative of the vegetation community and reflects the similar structure in the Coastal Sand Plains.

### Sampling and Data Collection

A vegetation map of the study area was created using land use/land cover (LULC) derived from ESA Sentinel-2 imagery at 10m resolution (Karra et. al. 2021). The underlying

deep learning model uses 6-bands of Sentinel-2 L2A surface reflectance data: visible blue, green, red, 2 shortwave infrared bands, and near-infrared (Karra et. al. 2021). The vegetation compositions I used in this study include - (1) woodland/shrubland composition which is categorized as any significant clustering of tall (~4.5 m or higher) dense vegetation; typically with closed or dense canopy such as wooded vegetation; clusters of dense tall vegetation within savannas, plantations, swamp, or mangroves (dense/tall vegetation with ephemeral water or canopy too thick to detect water underneath); (2) rangeland/wetland composition which is categorized as open areas covered in homogenous grasses with little to no taller vegetation; wild cereals and grasses with no obvious human plotting (i.e., not a plotted field); such as natural meadows and fields with sparse to no tree cover, open savanna with few to no trees or pastures. There may also be a mix of small clusters of plants or single plants dispersed on a landscape that shows exposed soil or rock; scrub-filled clearings within dense forests that are clearly not taller than trees; such as moderate to sparse cover of bushes, shrubs and tufts of grass, savannas with very sparse grasses, trees or other plants; these areas are of any type of vegetation with obvious intermixing of water throughout a majority of the year or seasonally; (3) sand composition is categorized as large areas of sand and deserts with no to little vegetation; fine rock or soil with very sparse to no vegetation for the entire year; such as sand dunes and desert, exposed fine rock or soil (Karra et al. 2021).


I deployed a total of 5 Wildlife Acoustic SM4 Bat FS Ultrasonic Recorders coupled with ultrasonic omnidirectional microphones and placed them within a vegetation composition from February-July 2022. The rangeland/wetland composition sampling took place during the months of February, May, and July at 15 different sites each per month for 5 consecutive days. The sand composition sampling took place during March with a total of 5 unique sites during 5

consecutive days. The woodland/shrubland composition sampling took place during April at 5 individual sites for 5 consecutive days once again (Figure 1). I chose the sampling sites based on the historically predominant areas of each vegetation composition, taking into consideration their accessibility. I determined the number of sites per sampling period (month) by the total number of available ultrasonic recorders. An assumed maximum distance for bat echolocations to be recorded in perfect conditions is 100 m or 0.1 km (SonoBat 2023). Therefore, all sampling locations are a minimum of 0.1 km from the ranch boundaries and a minimum of 0.1 km from a simultaneous sampling location to assume independence.

I reported bat calls throughout the high foraging hours starting 1 hour before sunset and ending 1 hour after sunrise based on geographic location. To ensure capture of the peak activity time of bats, an acoustic sample starting 4 hours post-sunset and 2 hours pre-sunrise is sufficient (Perks and Goodenough 2020). The number of acceptable sampling nights varies significantly depending on the season. Furthermore, 40-60% of species richness at a site for the majority of the year can be accomplished by sampling 2 to 5 nights (Skalak et al. 2012). Therefore, I ran the recorders for at least 5 consecutive nights.

I attached these external ultrasonic omnidirectional microphones at the top of a tall pole placed either in the center of open vegetation types (i.e., rangeland) or in an open pathway/clearing within dense vegetation (i.e., woodland) to prevent obstruction of sound by the surroundings. On the recorder, I turned on a 16k High Filter that applies a gradual filter that will attenuate signals starting below 16 kHz in frequency to help filter out human and weather noise, which provides a more definite recording of signals. The default sample rate was set at 256 kHz, which can record up to 128 kHz and capture all species of bats found in North America (Wildlife Acoustics 2023). Species analyzed for this study include - Mexican free-tailed bat (*Tadarida*

*brasiliensis*, Tabr), Evening bat (*Nycticeius humeralis*, Nyhu), Hoary bat (*Lasiurus cinereus*, Laci), Tri-colored bat (*Perimyotis subflavus*, Pesu), Eastern red (*Lasiurus borealis*, Labo), Northern yellow (*Lasiurus intermedius*, Lain), and Cave myotis (*Myotis velifer*, Myve).




**Figure 1. Sites on the ESR where acoustic data were collected for the following vegetation compositions: woodland/shrubland, rangeland/wetland, and sand.**

## Acoustic Data Analysis

I used SonoBat v. 4.4.5. as the primary method in analyzing the activity of various species of bats utilizing my study area (Figure 2). SonoBat allows me to identify species detected and their frequency of occurrence. SonoBat produces high-resolution sonograms of each call pulse to extract full call trends (SonoBat 2023). All frequency content at any point in time is encoded into each recording and an algorithm is run to conclude bat species. I manually confirmed any non-classified calls by referencing known calls published by Echolocation Call Characteristics of US Bats. To avoid overestimation, frequency of occurrence is necessary as observing the bats in real-time to determine the exact number of individuals was not feasible in this study. The same individual bat could have passed by the microphone multiple times and echolocation was documented multiple times appearing as more than one individual.

For each recorder/sampling location, 5 corresponding consecutive nights were analyzed for each sampling period and combined in final analysis for the 3 vegetation compositions. First, the sound files were attributed with meta-data (date, location, site, and time). I then analyze sequences collected by my acoustic recorders using the auto classifiers on the SonoBat Batch Scrubber. Using the batch of files, the software detects files without bat calls, such as those set off by noise, and relocates them to a separate folder to exclude them from further analysis. The default settings are selected: medium scrubbing, accept all calls except low-quality calls, accept some noise with tonal content, and include 5 kHz signals and above. Using the regional classifier “North American Regional Classifier”, species that are likely to be identified include Mexican Free-Tailed, Eastern Red, Cave Myotis, and Evening Bat (Fern et al. 2018). The remaining files from each package will include species-level identification of each bat pass, when applicable. The remaining files in each batch will indicate species-level identification for each bat pass, as

appropriate. Once the automated identification is complete, I manually inspect the sound files to confirm the correct identification of species, using reference calls found in the literature (Echolocation Call Characteristics of U.S. Bats 2011).



**Figure 2. SonoBat call analysis sonogram.**

### **Vegetation Selection Ratios**

I evaluated vegetation selection by studied bat species using a vegetation-selection ratio by all species and by individual species (Davis et al. 1994, Lopez et al. 2004). The total species vegetation-selection ratios were calculated by dividing the observed frequency of occurrence by the expected frequency of occurrence for all species in each vegetation composition. The observed frequency of occurrence was calculated for all species by use of SonoBat described in the previous section. The expected frequency of occurrence was calculated by first calculating the availability or proportion of a given vegetation composition ( $\text{Availability} = \text{area of vegetation} / \text{total area of ESR}$ ), and then multiplying this availability by the total frequency of occurrence for all species within all the vegetation compositions (Expected frequency of

occurrence=availability\*total frequency of occurrence throughout all vegetation compositions).

The vegetation-selection ratio ( $S$ ) was then calculated as  $S=U/A$ , where  $U$  is the observed frequency of occurrence and  $A$  is the expected frequency of occurrence when accounting for availability (Davis et al. 1994, Lopez et al. 2004).

Vegetation-selection ratios  $S>1$  suggests the bat frequency of occurrence was greater than the expected frequency of occurrence or showed a preference for, and  $S<1$  suggests the bat frequency of occurrence was less than the expected frequency of occurrence or showed avoidance of area (Davis et al. 1994, Lopez et al. 2004). Species specific vegetation-selection ratios were also calculated. This was achieved in a similar manner explained above with the exception of the “total frequency of occurrence throughout all vegetation compositions” replaced with the “total species-specific frequency of occurrence within the vegetation composition of interest.” Following graphs suggest overall preference or avoidance of a vegetation composition as well as which species of bats prefer or avoid each vegetation composition.

## RESULTS

### Frequency of Occurrence

I surveyed a total of 25 sites across ESR, with a total of 25 unique sampling nights analyzed throughout the entire sampling period. A total of 70,650 total wav files were recorded, with 20,518 being of adequate quality to use in the final analysis (29.04%). Seven species were used for the analysis with the Evening bat having the most frequency of occurrence overall vegetation compositions ( $n=14,417$ ) (Table 1). Other species detected included Eastern red bat ( $n=2,670$ ), Hoary bat ( $n=1,390$ ), Northern yellow bat ( $n=949$ ), Mexican free-tailed bat ( $n=573$ ), Tri-colored bat ( $n=476$ ), and Cave myotis bat ( $n=43$ ).


**Table 1. Scientific and common names for every species analyzed with total occurrence per species for the entirety of the study in decreasing order of occurrence.**

| Scientific Name              | Common Name             | Total Occurrence |
|------------------------------|-------------------------|------------------|
| <i>Nycticeius humeralis</i>  | Evening bat             | 14,417           |
| <i>Lasiurus borealis</i>     | Eastern red bat         | 2,670            |
| <i>Lasiurus cinereus</i>     | Hoary bat               | 1,390            |
| <i>Lasiurus intermedius</i>  | Northern yellow bat     | 949              |
| <i>Tadarida brasiliensis</i> | Mexican free-tailed bat | 573              |
| <i>Perimyotis subflavus</i>  | Tri-colored bat         | 476              |
| <i>Myotis velifer</i>        | Cave myotis bat         | 43               |

### Vegetation Selection

The rangeland/wetland had the highest availability of 81.79% of the area, followed by the woodland/shrubland composition of 12.57%, and finally, the sand composition had an availability of 4.76%. Included is an “other” category that accounts for areas I did not include in

my study such as developments (i.e., buildings) or roads that represent less than 1% of the area (0.89%) (Figure 3). A preference for an area would require the vegetation-selection ratio within this vegetation composition as  $S \geq 1$ .




**Figure 3. Vegetation availability by composition; rangeland/wetland (82%), woodland/shrubland (12%), sand (5%), other (1%).**

Results for the frequency of occurrence per species in each vegetation composition are found below (Table 2). The woodland/shrubland (WS) composition had the most total frequency of occurrence among the other 2 compositions studied ( $n=11,757$ ) consisting of over half (57.30%) of all activity in this vegetation composition alone. The rangeland/wetland (RW) composition had the second total frequency of occurrence ( $n=5,818$ ) accounting for 28.36% of all activity within this vegetation. The sand (SD) composition had the least total frequency of occurrence compared to the other vegetation compositions studied ( $n=2,943$ ), with only 4.34% of all activity.


**Table 2. Studied bat species with total occurrence per vegetation composition, woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

| Bat Species             | WS Occurrence | RW Occurrence | SD Occurrence |
|-------------------------|---------------|---------------|---------------|
| Eastern red bat         | 1,401         | 930           | 339           |
| Hoary bat               | 1,058         | 261           | 71            |
| Northern yellow bat     | 738           | 136           | 75            |
| Cave myotis bat         | 23            | 20            | 1             |
| Evening bat             | 8,087         | 4,106         | 2,224         |
| Tri-colored bat         | 215           | 250           | 11            |
| Mexican free-tailed bat | 235           | 115           | 223           |
| TOTAL                   | 11,757        | 5,818         | 2,943         |

The vegetation-selection ratios for each vegetation composition were calculated. The woodland/shrubland (WS) composition ( $S=4.56$ ) and sand (SD) composition ( $S=3.01$ ) both had total vegetation-selection ratios of  $S>1$ , showing a preference for these 2 areas. The rangeland/wetland (RW) composition ( $S=0.35$ ) had a total ratio of  $S<1$ , which shows avoidance of this vegetation composition overall (Figure 4). Although Figure 4 shows the comprehensive preference and avoidance for each vegetation composition, the unique bat species are also studied to demonstrate preference/avoidance of vegetation compositions for each individual species (Figure 5).




**Figure 4. Total vegetation-selection ratios ( $S$ ) ( $S > 1$  signifies a preference for the area,  $S < 1$  signifies an avoidance) for each vegetation composition, woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**



**Figure 5. Species specific selection ratios ( $S$ ) for each vegetation composition ( $S > 1$  signifies a preference for the area,  $S < 1$  signifies an avoidance), Eastern red bat (Labo), Hoary bat (Laci), Northern yellow bat (Lain), Cave myotis bat (Myve), Evening bat (Nyhu), Tri-colored bat (Pesu), Mexican free-tailed bat (Tabr), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

### *Eastern red bat*


I observed a total frequency of occurrence for the Eastern red bat throughout the entirety of the study of  $n=2,670$ . The Eastern red bats preferred the use of the woodland/shrubland vegetation ( $S=4.17$ ) as well as the sand composition ( $S=2.67$ , preferred). This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.43$ , avoided) (Figure 6).



**Figure 6. Eastern red bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

**Hoary bat**


I observed a total frequency of occurrence for the Hoary bat throughout the entirety of the study of  $n=1,390$ . The Hoary bats preferred the use of the woodland/shrubland vegetation ( $S=6.06$ ) as well as the sand composition ( $S=1.07$ , preferred). The sand composition preference was extremely close to what is expected frequency of occurrence within the area with a vegetation-selection ratio of almost  $S=1$ . This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.23$ , avoided) (Figure 7).



**Figure 7. Hoary bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

### *Northern yellow bat*


I observed a total frequency of occurrence for the Northern yellow bat throughout the entirety of the study of  $n=949$ . The Northern yellow bats preferred the use of the woodland/shrubland vegetation ( $S=6.19$ ) as well as the sand composition ( $S=1.66$ , preferred). This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.18$ , avoided) (Figure 8).



**Figure 8. Northern yellow bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

*Cave myotis bat*


I observed a total frequency of occurrence for the Cave myotis bat throughout the entirety of the study of  $n=43$ . The Cave myotis bats preferred the use of the woodland/shrubland vegetation ( $S=4.26$ , preferred). This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.57$ ) as well as the sand composition ( $S=0.004$ , avoided) (Figure 9).



**Figure 9. Cave myotis bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

### *Evening bat*


The Evening bats had the highest frequency of occurrence out of the species analyzed with  $n=14,417$  throughout the entirety of the study. The Evening bats preferred the use of the woodland/shrubland vegetation ( $S=4.46$ ) as well as the sand composition ( $S=3.24$ , preferred). This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.35$ , avoided) (Figure 10).



**Figure 10. Evening bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

***Tri-colored bat***


I observed a total frequency of occurrence for the Tri-colored bat throughout the entirety of the study of  $n=476$ . The Tri-colored bats preferred the use of the woodland/shrubland vegetation ( $S=3.59$ , preferred). This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.64$ ) as well as the sand composition ( $S=0.49$ , avoided) (Figure 11).



**Figure 11. Tri-colored bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

***Mexican free-tailed bat***

The Mexican free-tailed bat had the highest vegetation-selection ratio. I observed a total frequency of occurrence for the Mexican free-tailed bat throughout the entirety of the study of  $n=573$ . The Mexican free-tailed bats preferred the use of the woodland/shrubland vegetation ( $S=3.26$ ) as well as the highest preference ratio of the sand composition ( $S=8.18$ , preferred). This species seemed to have an avoidance of the rangeland/wetland vegetation composition ( $S=0.25$ , avoided) (Figure 12).



**Figure 12. Mexican free-tailed bat vegetation-selection ratios stratified by vegetation type ( $S>1$  signifies a preference for the area,  $S<1$  signifies an avoidance), woodland/shrubland (WS), rangeland/wetland (RW), and sand (SD).**

## DISCUSSION

Research on Texas bats is limited, particularly with fine resolution information on vegetation selection. In my study, I observed the greatest frequency of occurrence and highest habitat ratio for bats monitored to be in woodland/shrubland vegetation. Other studies (Fuentes-Montemayor et al. 2013 and Kurta et al. 2018) have reported that woodland areas play a critical role in providing bats with significant roosting and feeding opportunities, making them a frequently selected vegetation type for many bat species. For example, Fuentes-Montemayor (2013) showed higher bat activity in small patches of isolated woodlands rather than sparse woodlands. While the woodland/shrubland area accounted for only about 13% of the ESR study area, this area is likely highly desirable for the species of bats found in South Texas. The majority of the species are tree-roosting bats, likely preferring woody coverage for shelter. Previous studies report (Kalka et al. 2008, Kurta et al. 2018, Fuentes-Montemayor et al. 2013) that many North American species of bats depend on tree cavities, tree bark, and canopy foliage for roosting. The woodland/shrubland vegetation furnishes a shield against wind and predators, while simultaneously offering ample corridors for flight where flight activity has been recorded 3 times higher than in single tree areas (Barros et al. 2014, Kalda et al. 2015). The previous results align with a majority of studies within the subject and are corroborated by the outcomes of my own study. Conserving and managing woodlands in this area of Texas is recommended for maintaining bat communities.

The rangeland/wetland accounted for most of the ESR area (~82%); however, despite the extensive coverage of rangeland/wetlands in the ESR, bats avoided this vegetation. The wetlands were categorized together with the rangelands as they consisted of small ephemeral ponds situated atop the rangelands, therefore sharing the same spatial extent. One main reason for this

lack of acoustic data may be due to the time it takes for bats to use the wetlands as water sources. Bats take only milliseconds to intake a gulp of water as they drink “on the fly” within these wetland areas (Klopper et al. 2019). Therefore, the acoustic data given by the bats may not be significant enough for the acoustic monitors to collect and store that data to be used in analysis. Although these water sources are extremely important to bats to survive (Fukui et al. 2006, Flaquer et al. 2009, Lookingbill et al. 2010, Šuba et al. 2012, Mas et al. 2021) the use of acoustic monitoring of bat calls alone may not be the best possible way to analyze the utilization of wetlands (Anderson et al. 2021). Other methods such as visual surveys may be needed for more accurate results of usage (Klopper et al. 2019). Studies (Fukui et al. 2006, Flaquer et al. 2009, Lookingbill et al. 2010, Šuba et al. 2012, Mas et al. 2021) have clearly proved that wetlands are frequently utilized by bats and some bats even have a preference for these areas, but my study showed an avoidance of these wetlands. The previous studies (Fukui et al. 2006, Flaquer et al. 2009, Lookingbill et al. 2010, Šuba et al. 2012, Mas et al. 2021) did not occur on cattle production ranches, thus neglecting the influence of factors specific to those operations that could have impacted the outcome of my results. Active ranching practices for cattle production, like those active at this study site, use anthropogenic water sources such as stock tanks that may provide external water sources for bats and insects (Geluso et al. 2004, Morris et al. 2011, Trubitt et al. 2019). It is also possible that the avoidance in the area can be attributed to the insufficient tree coverage rather than the area itself as shown in the study by Trubitt in 2019. Different bat species prefer a wide variety of different percentages of woody coverage habitats (Trubitt et al. 2019). Two species of bats studied by both Trubitt and I included the Northern yellow bat and the Eastern red bat. For these two species, their activity increased as tree cover increased; but the Big brown bat responded negatively to tree cover, giving differing results that could explain the

contrasting findings (Trubitt et al. 2019). As for the rangeland portion of the composition, bats prey on a wide variety of insects, such as beetles, mosquitoes, and moths (Coutts et al. 1973, Bernard 2002, Kunz et al. 2011, Long and Kurta 2014, Woodward 2022). The prevalent prey species thrive in the expansive rangeland areas, readily accessible to bats and ensuring an abundant food supply (Coutts et al. 1973, Bernard 2002, Kunz et al. 2011, Long and Kurta 2014, Woodward 2022). Consequently, it is crucial to manage rangelands and wetlands in a manner that offers a conducive habitat for prey and other species, while also ensuring the availability of wetlands or alternative water sources to support their presence.

While the sampling methods used in this study did procure both species-specific and overall vegetation preference results, the main limitation of this study would be not sampling into the later months of the year and due to the number of monitors available and not sampling each vegetation composition simultaneously. Sampling each vegetation each month would allow for results to be compared temporally. Seasonality as well as abiotic factors such as temperature and humidity would possibly be factors that played a role in frequency of occurrence. For my study's species-specific results, the Evening bat species exhibited a prevailing prevalence compared to all other species. In 2015 and 2016, a study (Szewszak et al. 2022) was done at the ESR to collect bat call data for occupancy estimates (along with other species similar to species in this study) for ~5 sampling nights. The Evening bat had a resulting number of passes detected of 76 in 2015 and 114 in 2016 at 5 and 10 locations respectively (Szewszak et al. 2022). For both years, the Evening bat ranked second in terms of species with most passes detected at the ESR, while the Eastern red bat ranked first (Szewszak et al. 2022). For my study, the top 2 species of bats were flipped, with the Eastern red bat (13.01%) ranking second behind the Evening bat accounting for 70.27% of all the frequency of occurrence data (Table 4.1). One possible reason

for this may be due to the differences in sampling periods. The Szewszak et al. (2022) study concluded that the maximum detection sampling period for the Eastern red bat is late July - late August and early August - mid September for Evening bat. I surveyed much earlier in the season beginning in February and ending in July. By extending my study into September, I may have seen similar results to this previous study. Although, this does present the possibility that higher occurrences of the Evening bat may be seen in earlier months with higher pikes of the Eastern red bat in the later months. By utilizing a different study period and conducting additional analysis, it is possible to examine particular species in more detail or species of interest such as migratory species to explore which vegetation compositions these species prefer at this critical stage in life. Using information found from this study as a reference point is important to note for future studies within this area of Texas and will provide more comprehensive insights to inform future management decisions.

## MANAGEMENT IMPLICATIONS

Species diversity is vital for ecosystem stability, resilience, productivity, provision of ecological services, and the long-term survival of species themselves (Tilman 1996, Schwartz et al. 2000, Smet and Ward 2009, Oliver et al. 2015). Rangeland management practices should focus on landscape heterogeneity to create a foundation for ecological systems to support biodiversity (Christensen 1997, Wiens 1997, Bailey 1998). While woodland/shrubland areas may be regarded as paramount, scientists, biologists, and agriculturalists understand that species diversity cannot flourish solely based on a single factor. Just like any other species, bats need a diverse range of resources and amenities for their well-being and thriving. Woodland conservation should be of main concern in aspects of vegetation and roosting habitat conservation of bats. My study has shown the importance of woodland corridors for bats, but these areas also act as gateway access for other species (i.e., birds and small mammals) as well as cattle on rangelands (Trubitt et al. 2019). Although the wetlands area showed a general avoidance, recommendations for cattle tanks remain to provide an abundance of anthropogenic water sources for both species (Fuentes-Montemayor et al. 2013, Kurta et al. 2018, Trubitt et al. 2019). The conservation of native grasslands, woodlands, and water sources not only supports foraging bats but also facilitates the interconnectedness of operational cattle ranching operations, resulting in positive outcomes for multiple species.

## REFERENCES

- Agosta, S. J. 2002. Habitat use, diet, and roost selection by the big brown bat (*Eptesicus fuscus*) in North America: A case for conserving an abundant species. *Mammal Review* 32:179–198.
- Anderson, K., S. Ibrahim, C. Phuong, and Y. Triyono. 2021. Difference in preference for water sources in insectivorous bats at Sedgwick Reserve 5:6
- Bailey, D. W., B. Dumont, M. F. Wallisdevries. 1998. Utilization of heterogeneous grasslands by domestic herbivores: Theory to management. *Annals of Zootechnology* 47:321-333.
- Barros, M. S., D. A. Pessoa, and A. M. Rui. 2014. Habitat use and seasonal activity of insectivorous bats (Mammalia: Chiroptera) in the grasslands of southern Brazil. *Zoologia (Curitiba)* 31:153–161.
- Bernard, E. 2002. Diet, activity, and reproduction of bat species (Mammalia, Chiroptera) in Central Amazonia, Brazil. *Revista Brasileira de Zoologia* 19:173–188.
- Billeter, R., J. Liira, D. Bailey, R. Bugter, P. Arens, I. Augenstein, S. Aviron, J. Baudry, R. Bukacek, F. Burel, M. Cerny, G. De Blust, R. De Cock, T. Diekötter, H. Dietz, J. Dirksen, C. Dormann, W. Durka, M. Frenzel, R. Hamersky, F. Hendrickx, F. Herzog, S. Klotz, B. Koolstra, A. Lausch, D. Le Coeur, J. P. Maelfait, P. Opdam, M. Roubalova, A. Schermann, N. Schermann, T. Schmidt, O. Schweiger, M. J. M. Smulders, M. Speelmans, P. Simova, J. Verboom, W. K. R. E. Van Wingerden, M. Zobel, and P. J. Edwards. 2008. Indicators for biodiversity in agricultural landscapes: a pan-European study. *Journal of Applied Ecology* 45:141–150.

- Christensen, N. L. 1997. Managing for heterogeneity and complexity on dynamic landscapes. *The Ecological Basis for Conservation: Heterogeneity, Ecosystems, and Biodiversity* Pages 167-186.
- Campbell, A. 2022. Bat guano fertilizer: benefits and how to use. Dre Campbell Farm. <<https://drecampbell.com/bat-guano-fertilizer-benefits-how-to-use/>>. Accessed 13 February 2021.
- Cleland, E. E. 2011. Biodiversity and ecosystem stability. *Nature Education Knowledge* 3:14.
- Cooper, J., J. Masterson, J. Thapa, M. Patton, S. Limkhedawala, S. Parab, and S. Maharajan. 2021. State of community report Willacy County, Texas source: Texas historical commission.
- Coutts, R. A., M. B. Fenton, and E. Glen. 1973. Food intake by captive *Myotis lucifugus* and *Eptesicus fuscus* (Chiroptera: Vespertilionidae). *Journal of Mammalogy* 54:985–990.
- D’Antonio, C. M., and P. M. Vitousek. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23:63–87.
- Davis, A. J., Manly, B. F. J., McDonald, L. L., and D. L. Thomas. 1994. Resource selection by animals: statistical design and analysis for field studies. *Journal of Animal Ecology* 63:745.
- Echolocation Call Characteristics of U.S. Bats. 2011. Echolocation call characteristics of Eastern U.S. bats. SonoBat, Arcata, California.
- Fern, R. R., H. T. Davis, J. A. Baumgardt, M. L. Morrison, and T. A. Campbell. 2018. Summer activity patterns of four resident south Texas bat species. *Global Ecology and Conservation* 16:e00500.
- Fuentes-Montemayor, E., D. Goulson, L. Cavin, J. M. Wallace, and K. J. Park. 2013. Fragmented woodlands in agricultural landscapes: the influence of woodland character and

- landscape context on bats and their insect prey. *Agriculture, Ecosystems and Environment* 172:6–15.
- Gough, L., C. W. Osenberg, K. L. Gross, S. C. Gough, C. W. Gross, K. L. Collins, and C. W. Osenberg. 2000. Fertilization effects on species density and primary productivity in herbaceous plant communities. *Oikos* 89:428–439.
- Grashof-Bokdam, C. J., and F. Van Langevelde. 2005. Green veining: landscape determinants of biodiversity in European agricultural landscapes. *Landscape Ecology*. 20:417–439.
- Grime, J. 2001. Plant strategies, vegetation processes, and ecosystem properties. *Biological Conservation* 107:260–261.
- Wiens, J. A. 1997. The emerging role of patchiness in conservation biology. *The Ecological Basis for Conservation: Heterogeneity, Ecosystems, and Biodiversity* Pages 93-107.
- Kalda, O., R. Kalda, and J. Liira. 2015. Multi-scale ecology of insectivorous bats in agricultural landscapes. *Agriculture, Ecosystems and Environment* 199:105–113.
- Kalka, M. B., A. R. Smith, and E. K. V. Kalko. 2008. Bats limit arthropods and herbivory in a tropical forest. *Science* 320:71.
- Karra, K., C. Kontgis, Z. Statman-Weil, J. C. Mazzariello, M. Mathis, and S. P. Brumbly. 2021. Global land use/land cover with Sentinel-2 and deep learning. *IGARSS 2021-2021 IEEE International Geoscience and Remote Sensing Symposium*. Brussels, Belgium. Pages 4,704-4,707.
- Knight, S. 2020. Bats are important but some in Texas face uncertain future. *Tyler Morning Telegraph*. Tyler, Texas. M. Roberts Media. B5.
- Kloepper, L. N., A. M. Simmons, and J. A. Simmons. 2019. Echolocation while drinking: Pulse-timing strategies by high- and low-frequency FM bats. *PLOS ONE* 14:e0226114.

- Kunz, T. H., E. B. de Torrez, D. Bauer, T. Lobova, and T. H. Fleming. 2011. Ecosystem services provided by bats. *Annals of the New York Academy of Sciences* 1,223:1–38.
- Kurta, A., J. P. Hayes, and M. J. Lacki. 2018. Bats in forests: conservation and management. *Bats in Forests* 15:154-155.
- Loeb, S. C., and R. V Blakey. 2021. Bats and fire: a global review. *Fire Ecology* 17:29.
- Long, B. L., and A. Kurta. 2014. Activity and diet of bats in conventional versus organic apple orchards in southern Michigan. *Canadian Field-Naturalist* 1,282:158-164.
- Lopez, R. R., N. J. Silvy, N. R. Wilkins, P. A. Frank, M. J. Peterson, and N. M. Peterson. 2004. Habitat-use patterns of Florida key deer: implications of urban development. *Journal of Wildlife Management* 68:900-908.
- Marin, L. 2023. East Foundation Earns ARO Status. *The Land Report*. Featured Ranchland. Virginia Beach, Florida.
- Mittelbach, G. G., C. F. Steiner, S. M. Scheiner, K. L. Gross, H. L. Reynolds, R. B. Waide, M. R. Willig, S. I. Dodson, and L. Gough. 2001. What is the observed relationship between species richness and productivity? *Ecology* 82:2381.
- Perks, S. J., and A. E. Goodenough. 2020. Abiotic and spatiotemporal factors affect activity of European bat species and have implications for detectability for acoustic surveys. *Wildlife Biology* 2020.
- PRISM Climate Group. 2023. PRISM Climate Group at Oregon State University.
- Richardson, D. M., P. Py S Ek, M. Rejmánek, M. G. Barbour, F. D. Panetta, and C. J. West. 2000. Naturalization and invasion of alien plants: concepts and definitions. *Wiley Online Library* 6:93–107.

- Robinson, R. A., and W. J. Sutherland. 2002. Post-war changes in arable farming and biodiversity in Great Britain. *Journal of Applied Ecology* 39:157–176.
- Russo, D., L. Bosso, and L. Ancillotto. 2018. Novel perspectives on bat insectivory highlight the value of this ecosystem service in farmland: Research frontiers and management implications. *Agriculture Ecosystems and Environment* 266:31–38.
- Russo, D., and G. Jones. 2002. Identification of twenty-two bat species (Mammalia: Chiroptera) from Italy by analysis of time-expanded recordings of echolocation calls. *The Zoological Society of London* 91–103.
- Schnitzler, H. U., and E. Kalko. 2001. *Echolocation by insect-eating bats*. Oxford Academic. *BioScience* 51:557–569.
- Sekercioglu, C. H. 2006. Increasing awareness of avian ecological function. *Trends in Ecology and Evolution* 21:464–471.
- Simberloff, D., J. L. Martin, P. Genovesi, V. Maris, D. A. Wardle, J. Aronson, F. Courchamp, B. Galil, E. García-Berthou, M. Pascal, P. Pyšek, R. Sousa, E. Tabacchi, and M. Vilà. 2013. Impacts of biological invasions: what’s what and the way forward. *Trends in Ecology and Evolution* 28:58–66.
- Skalak, S. L., R. E. Sherwin, and R. M. Brigham. 2012. Sampling period, size, and duration influence measures of bat species richness from acoustic surveys. *Methods in Ecology and Evolution* 3:490–502.
- SonoBat. 2023. SonoBat – Software for Bat Call Analysis. SonoBat, Arcata, California. <<https://sonobat.com/>>. Accessed 5 January 2022.

- Steel, Z. L., B. Campos, W. F. Frick, R. Burnett, and H. D. Safford. 2019. The effects of wildfire severity and pyrodiversity on bat occupancy and diversity in fire-suppressed forests. *Scientific Reports* 9:16,300.
- Szewszak, J. M., T. A. Campbell, J. A. Baumgardt, M. L. Morrison, L. A. Brennan, H. T. Davis, R. R. Fern. 2022. Monitoring occupancy of bats with acoustic data: power and monitoring occupancy of bats with acoustic data: power and sample size recommendations sample size recommendations. *Western North American Naturalist*. 82:36–49.
- Texas Parks and Wildlife Department. TEAM: Texas Ecosystem Analytical Mapper. Austin, Texas. <<https://tpwd.texas.gov/gis/team/>>. Accessed 10 March 2022.
- Torquetti, C. G., A. T. B. Guimarães, and B. Soto-Blanco. 2021. Exposure to pesticides in bats. *The Science of the Total Environment* 755:1,425,009.
- Wied, J. P., H. L. Perotto-Baldivieso, A. A. T. Conkey, L. A. Brennan, and J. M. Mata. 2020. Invasive grasses in South Texas rangelands: historical perspectives and future directions. *Invasive Plant Science and Management* 13:41–58.
- Wildlife Acoustics. 2023. Song Meter SM4BAT-FS Ultrasonic Bat Detector. Wildlife Acoustics.
- Woodward, L. 2022. Bat Conservation Texas Style. Texas Wildlife Association. New Braunfels, Texas. <<https://tpwd.texas.gov/huntwild/wild/species/bats/bat-watching-sites/>>. Accessed 7 February 2022.